arXiv:1301.6071v1 [math.PR] 25 Jan 2013

A local CLT for convolution equations with an application to weakly self-avoiding random walks[*]

Luca Avena[†] Erwin Bolthausen[‡] Christine Ritzmann
University of Zürich

Abstract

We prove error bounds in a central limit theorem for solutions of certain convolution equations. The main motivation for investigating these equations stems from applications to lace expansions, in particular to weakly self-avoiding random walks in high dimensions. As an application we treat such self-avoiding walks in continuous space. The bounds obtained are sharper than the ones obtained by other methods.

1 Introduction

Let ϕ be the standard normal density in $\mathbb{R}^d$, $\mathbf{B} = \{B_k\}_{k\geq 1}$ be a sequence of rotationally invariant integrable functions, and $\lambda > 0$ a (small) parameter. Define recursively

$$\begin{aligned} C_0 &= \delta_0, \qquad (1.1)\\ C_n &= C_{n-1} * \phi + \lambda \sum_{k=1}^{n} c_k B_k * C_{n-k},\ n \geq 1, \end{aligned}$$

where

$$c_n \overset{\text{def}}{=} \int C_n(x)\, dx.$$

δ_0 denotes the Dirac "function". All our (signed) distributions will have densities, except those at index 0.

As written above, the sequence $\mathbf{C} = \{C_n\}_{n\geq 0}$ is not quite recursively defined as the right hand side contains the summand $c_n B_n$. The sequence $\{c_n\}$ itself satisfies

$$\begin{aligned} c_0 &= 1, \qquad (1.2)\\ c_n &= c_{n-1} + \lambda \sum_{k=1}^{n} c_k b_k c_{n-k},\ n \geq 1, \end{aligned}$$

---

[*] AMS 2000 classification: 60K35, 60F05. Key words and phrases: central limit theorem, convolution equations, self-avoiding random walks.

[†] supported by the Swiss National Science Foundation under contract 138141, and by the Forschungskredit of the University of Zürich.

[‡] supported by the Swiss National Science Foundation under contract 138141, and by the Humboldt Foundation.

where $b_k = \int B_k(x)\,dx$. Therefore, if $\lambda |b_n| < 1$ for all n, these equations define the sequence $\{c_n\}$ uniquely, and then, also $\mathbf{C}$ is well-defined. We will always assume that we are in this situation.

The main assumption is a decay property of the B_n for large n. We will also assume Gaussian decay properties in space which are natural for the applications to self-avoiding walks. The method we present here can probably be adapted to treat situations with less severe decay assumptions in space, but we have not worked that out.

Our main interest is to prove a local central limit theorem for C_n/c_n under appropriate conditions on $\mathbf{B}$ and λ. Of course, the parameter λ can be incorporated into $\mathbf{B}$. However, the approach we follow is purely perturbative. We will give conditions on $\mathbf{B}$, and then state that if in addition λ is small enough a CLT holds.

We use ϕ as the convoluting factor in (1.1) with C_{n-1}, so that for $\lambda = 0$, C_n simply is the normal density with covariance matrix $n \times$ identity. At the expense of a few complications, we could also investigate the case where the first summand in (1.1) is $C_{n-1} * S$ with a rotationally invariant density S. We however feel that this generalization would somehow obscure the main line of the argument. To step out from the rotationally invariant case however leads to new, complicated, and interesting problems which will be presented elsewhere.

The main motivation for our investigation of the equations comes from weakly self-avoiding random walks, as first investigated by Brydges and Spencer in the seminal paper [2]. Their results are for random walks on the d-dimensional lattice $\mathbb{Z}^d,\ d \geq 5$. In contrast, we investigate weakly self-avoiding random walks on $\mathbb{R}^d$ with standard normal increments. The model has two parameters $\lambda, \rho > 0$, ρ being the range of the interaction, and λ the strength. We set $\mathbb{I}_\rho(x) \overset{\mathrm{def}}{=} 1_{\{|x| \leq \rho\}}$, and if $\mathbf{x} = (x_1, \ldots, x_n) \in \left(\mathbb{R}^d\right)^n$, and $0 \leq i < j \leq n$, we set $U^\rho_{ij}(\mathbf{x}) \overset{\mathrm{def}}{=} \mathbb{I}_\rho(x_j - x_i)$, where $x_0 = 0$. Then, for $0 \leq \lambda \leq 1$, define the probability measure $P_{n,\lambda,\rho}$ on $\left(\mathbb{R}^d\right)^n$ by its density with respect to Lebesgue measure:

$$p_{n,\lambda,\rho}(\mathbf{x}) = \frac{1}{Z_{n,\lambda,\rho}} K_{\lambda,\rho}[0,n](\mathbf{x})\, \Phi[0,n](\mathbf{x}),$$

where

$$K_{\lambda,\rho}[a,b](\mathbf{x}) \overset{\mathrm{def}}{=} \prod_{a \leq i < j \leq b} \left(1 - \lambda U^\rho_{ij}(\mathbf{x})\right), \tag{1.3}$$

$$\Phi[a,b](\mathbf{x}) \overset{\mathrm{def}}{=} \prod_{i=a+1}^{b} \phi(x_i - x_{i-1}). \tag{1.4}$$

$Z_{n,\lambda,\rho}$ is the usual partition function, i.e. the norming factor which makes $p_{n,\beta,\rho}$ into a probability density. The main interest is to prove a central limit theorem for this measure, in the simplest case for the last marginal measure. It is convenient to consider first the unnormalized kernel $C^{\mathrm{SAW}}_n(x)$, $x \in \mathbb{R}^d$, which is defined to be the last marginal density of $Z_{n,\beta}p_{n,\beta,\rho}(\mathbf{x})$, i.e.

$$C_n^{\mathrm{SAW}}(x_n) = \int K_{\lambda,\rho}[0,n](\mathbf{x})\,\Phi[0,n](\mathbf{x}) \prod_{i=1}^{n-1} dx_i. \tag{1.5}$$

By a lace expansion, the C_n^{SAW} satisfy an equation

$$C_n^{\mathrm{SAW}} = C_{n-1}^{\mathrm{SAW}} * \phi + \sum_{k=1}^{n} \Pi_k * C_{n-k}^{\mathrm{SAW}}, \tag{1.6}$$

where the kernels Π_k describe the interactions through the weak self-avoidance. The Π_k are complicated functions and are hard to evaluate precisely. However, one crucial property is that the leading order decay is the same as that of the C_k^{SAW}. It therefore looks natural to write $\Pi_k = \lambda c_k^{\mathrm{SAW}} B_k$, and one seeks for condition on the B_k ensuring a CLT for solutions of (1.1). We can then apply the central limit theorem obtained for (1.1), Theorem 2.2, provided we can check the conditions on the sequence $\mathbf{B}$. The theorem we obtain as a corollary of Theorem 2.2 is

Theorem 1.1
For $d \geq 5$, $\rho \in (0,1]$, and $0 < \varepsilon \leq \frac{1}{100}$, there exists $\lambda_0(d,\varepsilon) > 0$ such that for $\lambda \in (0,\lambda_0]$ there exist a parameter $\delta(d,\rho,\lambda) > 0$ and a constant $K(d,\varepsilon,\lambda) > 0$ such that for all $n \in \mathbb{N}$

$$\left|\frac{C_n(x)}{c_n} - \phi_{n\delta}(x)\right| \leq K\left[r_n \phi_{n\delta(1+\varepsilon)}(x) + n^{-d/2} \sum_{j=1}^{\lceil n/2 \rceil} j\phi_{j\delta(1+\varepsilon)}(x)\right]$$

with

$$r_n = \begin{cases} n^{-1/2} & \text{for } d = 5 \\ n^{-1}\log n & \text{for } d = 6 \\ n^{-1} & \text{for } d \geq 7 \end{cases}. \tag{1.7}$$

Remark 1.2

a) The bound leads to $\|C_n/c_n - \phi_{n\delta}\|_1 = O(r_n)$.

b) The theorem does not give a local CLT as at $x = 0$ both $\phi_{n\delta}(0)$ and the bound are of order $n^{-d/2}$. A moments reflection however reveals that there cannot be a local CLT as the starting point keeps to have a noticeable influence on $C_n(x)/c_n$ for points x at distance of order 1 from the origin. However, our bound proves

$$\lim_{r\to\infty} \limsup_{n\to\infty} \sup_{x:|x|\geq r} n^{d/2} \left|\frac{C_n(x)}{c_n} - \phi_{n\delta}(x)\right| = 0.$$

So the result comes as close as possible to a local CLT.

c) The summation up to $\lceil n/2 \rceil$ is somewhat arbitrary, and can be replaced by $\lceil \alpha n \rceil$ for any $\alpha \in (0,1)$, adapting K. In fact, for $0 < \alpha < 1$

$$n^{-d/2} \sum_{j=\lceil \alpha n \rceil}^{n} j\phi_{j\delta(1+\varepsilon)}(x) \leq K(\alpha) r_n \phi_{n\delta(1+\varepsilon)}(x)$$

We have chosen $\alpha = 1/2$ for convenience. The second summand however is important and it takes care of the failure of the local CLT for x near the origin.

Self-avoiding random walks in continuous space have never been investigated, to our knowledge. For our approach, continous space is actually more convenient than the lattice. The method we use is the one developed originally in the thesis of Christine Ritzmann [3], [1], but there are a number of improvements, generalizations, and simplifications presented here.

There is a huge literature on the lace expansions and applications to many types of models. For a survey of the state of art around 2006, see [4], but there were still many developments since then. The emphasize here is to present an elementary and completely self-contained proof of a sharp CLT for solutions of (1.1), together with the perhaps simplest possible application. No knowledge of earlier versions of lace expansions or [3] are assumed.

The basic analysis of (1.1) here is independent of the application to self-avoiding walks, and goes through in all dimensions.

2 Main results

We fix some notations: $\mathbb{N}$ is the set of natural numbers $\{1, 2, \ldots,\}$ and $\mathbb{N}_0 \overset{\text{def}}{=} \mathbb{N} \cup \{0\}$. For $t > 0$, ϕ_t is the centered normal density in $\mathbb{R}^d$ with covariance matrix $t \times$ identity. We write ϕ for ϕ_1. For $a \in \mathbb{R}$ we write $\mathbf{a}$ for the sequence identical to a.

We typically drop $*$ for the convolution, so we just write AB instead of $A * B$ for two integrable functions A, B.

We write $C_*(\mathbb{R}^d)$ for the set of continuous, integrable functions $f : \mathbb{R}^d \to \mathbb{R}$, vanishing at ∞, which are of the form $f(x) = f_0(|x|)$ for some continuous function $f_0 : [0, \infty) \to \mathbb{R}$. We also write $C_*^+(\mathbb{R}^d)$ for the strictly positive ones. Occasionally, we simply write C_* and C_*^+.

Here are the conditions we need for $\mathbf{B}$:

Condition 2.1
We assume that the functions $B_m \in C_*(\mathbb{R}^d)$ are dominated in absolute value by functions $\Gamma_m \in C_*^+(\mathbb{R}^d)$ which satisfy the following conditions:

B1 There exist numbers $\chi_n(s) > 0,\ 1 \leq s \leq n$, satisfying $\chi_n(s) = \chi_n(n-s)$, and for some constant K_1

$$\sum_{s=1}^{n-1} (s \wedge (n-s)) \chi_n(s) \leq K_1,\ \forall n, \tag{2.1}$$

such that

$$\Gamma_m \Gamma_n \leq \chi_{m+n}(m)\, \Gamma_{n+m}, \tag{2.2}$$

B2 *There exists a constant* $K_2 > 0$ *such that for* $t \leq s \leq 2t$ *one has*

$$\Gamma_s \leq K_2 \Gamma_{2t} \tag{2.3}$$

B3 *There exists* $K_3 > 0$ *such that for* $m \leq t,\ m \in \mathbb{N},\ t \in \mathbb{R}^+,\ k = 0,1,2,$ *one has*

$$\int \phi_t(x-y)\, |y|^{2k}\, \Gamma_m(y)\, dy \leq K_3 \gamma^{(k)}(m)\, \phi_{t+m}(x), \tag{2.4}$$

where

$$\gamma^{(k)}(m) \overset{\text{def}}{=} \int |y|^{2k}\, \Gamma_m(y)\, dy.$$

B4

$$K_4 \overset{\text{def}}{=} \sum_n n\gamma^{(0)}(n) < \infty,\ K_5 \overset{\text{def}}{=} \sum_n \gamma^{(1)}(n) < \infty,\ K_6 \overset{\text{def}}{=} \sum_n n^{-1}\gamma^{(2)}(n) < \infty. \tag{2.5}$$

A simple example where the conditions are satisfied is $\Gamma_n = n^{-a}\phi_{n/2},\ a > 2$, but unfortunately, the application to self-avoiding walks needs a slightly more complicated choice, as will be discussed later.

We occasionally use γ_n instead of $\gamma^{(0)}(n)$.

We remark that under the above condition, one has for

$$b_n \overset{\text{def}}{=} \int B_n(x)\, dx$$

the estimate

$$|b_n| \leq \gamma_n$$

with

$$\gamma_m \gamma_n \leq \chi_{m+n}(m)\, \gamma_{n+m}. \tag{2.6}$$

We formulate our basic result:

We fix an arbitrary positive $\varepsilon > 0$, and write

$$\psi_n \overset{\text{def}}{=} \phi_{n\delta(1+\varepsilon)}, \tag{2.7}$$

with δ defined below in (2.16).

In the rest of this section, we will use L as a positive constant, not necessarily the same at different occurences, which may depend on $d, \varepsilon, K_1 - K_6$, but not on n, λ.

Let

$$\zeta_1(n) \overset{\text{def}}{=} 1 + \sum_{i=0}^{2} \sum_{m=1}^{n} m^{2-i} \gamma^{(i)}(m) \tag{2.8}$$

$$\zeta_2(n) \overset{\text{def}}{=} \sum_{m=n}^{\infty} \left(\gamma^{(1)}(m) + m\gamma^{(0)}(m) \right),$$

$$\overline{\zeta}(n) \overset{\text{def}}{=} n^{-2} \sum_{j=1}^{n} \zeta_1(j) + n^{-1} \sum_{j=1}^{n} \zeta_2(j).$$

Because of (2.3) and (2.5) we have

$$\lim_{n\to\infty} \overline{\zeta}(n) = 0, \ \sum_n n^{-1}\overline{\zeta}(n) < \infty, \ \overline{\zeta}(m) \le \overline{\zeta}(2n) \ \text{ for } n \le m \le 2n. \tag{2.9}$$

Theorem 2.2
Assume Condition 2.1. Then, if λ is small enough (depending on d, ε and K_1-K_6),the following estimates holds

$$|C_n(x)/c_n - \phi_{n\delta}(x)| \le L\lambda \left[\sum_{s=1}^{[n/2]} s\left(\psi_s \Gamma_{n-s}\right)(x) + \overline{\zeta}(n)\psi_n(x) \right], \tag{2.10}$$

where $\delta = \delta(\mathbf{B}, \lambda) > 0$ is defined in (2.16).

In the example $\Gamma_n(x) = n^{-a}\phi_{n/2}(x)$, $2 < a < 3$, one has $\zeta_1(n) = \text{const} \times n^{3-a}$, $\zeta_2(n) = \text{const} \times n^{2-a}$, and therefore $\overline{\zeta}(n) = \text{const} \times n^{2-a}$, and therefore

$$|C_n(x)/c_n - \phi_{n\delta}(x)| \le L n^{2-a}\psi_n$$

giving a local CLT with error estimate. For $a > 3$, we get

$$|C_n(x)/c_n - \phi_{n\delta}(x)| \le L n^{-1}\psi_n.$$

As remarked above, this Γ_n cannot work for the application to self-avoiding walks, and in fact, a pure local CLT is not possible in that case.

A first question we address is about the behavior of the sequence $\{c_n\}$:

Proposition 2.3
Assume Condition 2.1 and let $\mathbf{c}$ be the sequence defined by (1.2). Then if λ is small enough the following holds:

a) There exists a unique $\mu > 0$ such that $\alpha \overset{\text{def}}{=} \lim_{n\to\infty} \mu^{-n} c_n$ exists in $(0, \infty)$.

b) Writing $a_n \overset{\text{def}}{=} \mu^{-n} c_n$, one has

$$|a_{n+1} - a_n| < L\lambda\overline{\gamma}_n \overset{\text{def}}{=} L\lambda \sum\nolimits_{j=n}^{\infty} \gamma_j. \tag{2.11}$$

c)

$$\mu^{-1} = 1 - \lambda \sum\nolimits_{k=1}^{\infty} a_k b_k. \tag{2.12}$$

Remark 2.4

a) Plugging the expression (2.12) into (1.2), we see that $\mathbf{a} = \{a_n\}_{n\in\mathbb{N}_0}$ satisfies $a_0 = 1$, and

$$a_n = a_{n-1} - \lambda a_{n-1} \textstyle\sum_{k=n+1}^{\infty} a_k b_k + \lambda \displaystyle\sum_{k=1}^{n} a_k b_k \left(a_{n-k} - a_{n-1}\right), \ n \geq 1. \tag{2.13}$$

b) From (2.11) we get

$$|a_n - \alpha| \leq L\lambda \sum_{k=n}^{\infty} k\gamma_k. \tag{2.14}$$

Proof of Proposition 2.3. Let $l_1(\mathbb{N})$ be the Banach space of absolutely summable sequences $\mathbf{x} = \{x_n\}_{n\in\mathbb{N}}$, and $l_\gamma(\mathbb{N})$ be the set of sequences with $\|\mathbf{x}\|_\gamma \overset{\text{def}}{=} \sup_n \overline{\gamma}_n^{-1} |x_n| < \infty$. $\left(l_\gamma(\mathbb{N}), \|\cdot\|_\gamma\right)$ is a Banach space, too, and by (2.5), $l_\gamma \subset l_1$, and the embedding is continuous. The linear map $s : l_1(\mathbb{N}) \to l_\infty(\mathbb{N}_0)$ is defined by $s(\mathbf{x})_0 = 0$, and $s(\mathbf{x})_n \overset{\text{def}}{=} \sum_{j=1}^{n} x_j,\ n \geq 1$. Evidently, $\|s(\mathbf{x})\|_\infty \leq \|\mathbf{x}\|_1 \leq L\|\mathbf{x}\|_\gamma$. We also define the affine mapping $S : l_1 \to l_\infty$ by $S(\mathbf{x}) \overset{\text{def}}{=} \mathbf{1} + s(\mathbf{x})$. We define two mappings ψ_1, ψ_2 from $l_1(\mathbb{N})$ to the set of sequences with index set $\mathbb{N}$. $\psi_1(\mathbf{x})_1 = \psi_2(\mathbf{x})_1 = 0$, and

$$\psi_1(\mathbf{x})_n \overset{\text{def}}{=} S(\mathbf{x})_{n-1} \sum_{k=n+1}^{\infty} b_k S(\mathbf{x})_k,$$

$$\psi_2(\mathbf{x})_n \overset{\text{def}}{=} \sum_{k=2}^{n} S(\mathbf{x})_k b_k \left[s(\mathbf{x})_{n-k} - s(\mathbf{x})_{n-1}\right]$$

for $n \geq 2$. Finally we set $\psi \overset{\text{def}}{=} \lambda\psi_1 + \lambda\psi_2$. Remark first that

$$\psi(\mathbf{0})_n = \lambda\psi_1(\mathbf{0})_n = \lambda \sum_{k=n+1}^{\infty} b_k,$$

so that $\|\psi(\mathbf{0})\|_\gamma \leq L\lambda$, by (2.6).

$$\begin{aligned}|\psi_1(\mathbf{x})_n - \psi_1(\mathbf{y})_n| &\leq \|s(\mathbf{x}) - s(\mathbf{y})\|_\infty \left[\sum_{k=n+1}^{\infty} |b_k S(\mathbf{x})_k| + \left|S(\mathbf{y})_{n-1}\right| \sum_{k=n+1}^{\infty} |b_k|\right] \\ &\leq L\|\mathbf{x} - \mathbf{y}\|_\gamma \left[2 + L\|\mathbf{x}\|_\gamma + L\|\mathbf{y}\|_\gamma\right] \sum_{k=n+1}^{\infty} \gamma_k\end{aligned}$$

$$\|\psi_2(\mathbf{x}) - \psi_2(\mathbf{y})\|_\gamma \leq L\|\mathbf{x} - \mathbf{y}\|_\gamma \left(1 + \|\mathbf{x}\|_\gamma + \|\mathbf{y}\|_\gamma\right).$$

Similarly, for $n \geq 2$, by resummation

$$\begin{aligned} \psi_1(\mathbf{x})_n - \psi_1(\mathbf{y})_n &= \sum_{j=1}^{n-1} x_j \sum_{k=n-j+1}^{n} \left(S(\mathbf{y})_k - S(\mathbf{x})_k\right) b_k \\ &\quad + \sum_{j=1}^{n-1} (y_j - x_j) \sum_{k=n-j+1}^{n} S(\mathbf{y})_k \, b_k. \end{aligned} \tag{2.15}$$

In the first summand, we estimate $\left|S(\mathbf{x})_k - S(\mathbf{y})_k\right|$ by $L \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma$, so we get for this part an estimate

$$\leq L \left\|\mathbf{x}\right\|_\gamma \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma \sum_{j=1}^{n-1} \sum_{t=j}^{\infty} \gamma_t \sum_{k=n-j+1}^{n} \gamma_k.$$

$$\begin{aligned} \sum_{j=1}^{n-1} \sum_{t=j}^{\infty} \gamma_t \sum_{k=n-j+1}^{n} \gamma_k &\leq \sum_{j=1}^{n-1} \sum_{t=j}^{\infty} \sum_{k=n-j+1}^{n} \chi_{t+k}(t) \, \gamma_{t+k} \\ &\leq \sum_{s=n+1}^{\infty} \gamma_s \sum_{t=1}^{s-1} \delta(s,t) \, \chi_s(t), \end{aligned}$$

where we have used (2.6), and where $\delta(s,t)$ is the number of indices j satisfying $1 \leq j \leq n-1$, $t \geq j$, $n-j+1 \leq s-t \leq n$, so that $\delta(s,t) \leq t \wedge (s-t)$, and using (2.1), we get for the first summand of (2.15) an estimate $\leq L \left\|\mathbf{x}\right\|_\gamma \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma \overline{\gamma}(n)$. In a similar way, we get for the second summand an estimate $\leq L \left(1 + \left\|\mathbf{y}\right\|_\gamma\right) \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma \overline{\gamma}(n)$ and therefore

$$\left\|\psi_2(\mathbf{x}) - \psi_2(\mathbf{y})\right\|_\gamma \leq L \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma \left(1 + \left\|\mathbf{x}\right\|_\gamma + \left\|\mathbf{y}\right\|_\gamma\right),$$

leading to

$$\left\|\psi(\mathbf{x}) - \psi(\mathbf{y})\right\|_\gamma \leq L\lambda \left\|\mathbf{x}-\mathbf{y}\right\|_\gamma \left(1 + \left\|\mathbf{x}\right\|_\gamma + \left\|\mathbf{y}\right\|_\gamma\right).$$

From that and $\psi(\mathbf{0}) \in l_\gamma$, it follows that ψ maps l_γ continuously into itself, and furthermore, if λ is small enough, by the Banach fixed point theorem, the iterates $\psi^n(\mathbf{0})$ converge in l_γ to an element ξ with $\left\|\xi\right\|_\gamma \leq L\lambda$ which is a fixed point of ψ.

If we write

$$\eta \overset{\text{def}}{=} S(\xi), \ \varpi \overset{\text{def}}{=} \left(1 - \lambda \sum\nolimits_{k=1}^{\infty} \eta_k b_k\right)^{-1},$$

then it is readily checked, using the fact that ξ is a fixed point of ψ, that the sequence $\{\eta_n \varpi^n\}$ satisfies (1.2), and therefore it *is* this sequence. So it follows that $\varpi = \mu$, and $\mu^{-n} c_n$ satisfies the properties listed in a)-c). ■

Let $\mathbf{f} = \{f_n\}$ be a sequence of functions in C_*^+ which satisfy $\lim_{n\to\infty} \sup_x f_n(x) = 0$. For any sequence $\mathbf{g} = \{g_n\}$, $g_n \in C_*$ define

$$\left\|\mathbf{g}\right\|_{\mathbf{f}} \overset{\text{def}}{=} \sup_n \sup_{x \in \mathbb{R}^d} \frac{\left|g_n(x)\right|}{f_n(x)},$$

and write $B_{\mathbf{f}} \overset{\text{def}}{=} \{\mathbf{g} : \|\mathbf{g}\|_{\mathbf{f}} < \infty\}$ which equipped with $\|\cdot\|_{\mathbf{f}}$ is a Banach space.

As $B_m \in C_*$, the "covariance" matrix satisfies

$$\int x^T x B_m(x)\, dx = \overline{b}_m I_d,$$

for some $\overline{b}_m \in \mathbb{R}$ (possibly negative), I_d being the $d \times d$ unit matrix. Evidently, $\left|\overline{b}_m\right| \leq \gamma^{(1)}(m)$, and by Conditions 2.1 (2.5), the following number is well defined (for small enough λ)

$$\delta \overset{\text{def}}{=} \frac{\mu^{-1} + \lambda \sum_{m=1}^{\infty} a_m \overline{b}_m}{\mu^{-1} + \lambda \sum_{m=1}^{\infty} m a_m b_m}. \tag{2.16}$$

By choosing $\lambda > 0$ small enough, we can achieve that

$$|1 - \delta| \leq L\lambda,\ |1 - \mu| \leq L\lambda. \tag{2.17}$$

and also

$$1/2 \leq a_n \leq 3/2,\ \forall n,$$

which we assume henceforward.

Let $\mathbf{C}$ be the solution of (1.1). We put $A_n \overset{\text{def}}{=} C_n \mu^{-n}$ where μ is given by (2.12). This sequence satisfies $A_0 = \delta_0$ and

$$A_n = \mu^{-1} A_{n-1} \phi + \lambda \sum_{k=1}^{n} a_k B_k A_{n-k}. \tag{2.18}$$

Then $a_n = \int A_n(x)\, dx$, and $A_n / a_n = C_n / c_n$.

Proof of Theorem 2.2. Evidently, the statement is the same (given Proposition 2.3) as to bound $|A_n(x) - a_n \phi_{n\delta}(x)|$ in the same way.

We define an operator Ψ on sequences of functions $\mathbf{G} = \{G_n\}_{n \geq 0}$, $G_n \in C_*$, $\Psi(\mathbf{G})_0 \overset{\text{def}}{=} G_0$, and for $n \geq 1$

$$\Psi(\mathbf{G})_n \overset{\text{def}}{=} a_n \phi_{n\delta} G_0 - \sum_{j=1}^{n} G_{n-j} \Delta(j, j),$$

with

$$\Delta(k, j) \overset{\text{def}}{=} a_j \phi_{k\delta} - \mu^{-1} a_{j-1} \phi_{(k-1)\delta+1} - \lambda \sum_{m=1}^{j} a_m a_{j-m} B_m \phi_{(k-m)\delta} \tag{2.19}$$

for $k \geq j$. A simple resummation gives

$$\Psi(\mathbf{G})_n = G_n - \sum_{j=1}^{n} a_{n-j} \phi_{(n-j)\delta} \left[G_j - \mu^{-1} \phi G_{j-1} - \lambda \sum_{m=1}^{j} a_m B_m G_{j-m} \right].$$

From that it follows that if $\mathbf{A}$ satisfies $A_0 = \delta_0$ and (2.18), then $\Psi(\mathbf{A}) = \mathbf{A}$, and vice versa: If $A_0 = \delta_0$, and $\mathbf{A}$ satisfies the fixed point equation, then (2.18) follows by induction on n.

We consider the Banach space $(B_{\mathbf{f}}, \|\cdot\|_{\mathbf{f}})$ where

$$f_n \overset{\text{def}}{=} \sum_{s=1}^{[n/2]} s\psi_s \Gamma_{n-s} + \overline{\zeta}(n)\,\psi_n. \tag{2.20}$$

If $\mathbf{E}$ is the sequence $\{a_n \phi_{n\delta}\}$ then $\Psi(\mathbf{E})_n = E_n - \sum_{j=1}^n a_{n-j}\Delta(n,j)$. From Lemma 3.1 (3.4), we see that $\Psi(\mathbf{E}) - \mathbf{E} \in B_{\mathbf{f}}$ with $\|\Psi(\mathbf{E}) - \mathbf{E}\|_{\mathbf{f}} \le L\lambda$. Furthermore, if $\mathbf{G} \in B_{\mathbf{f}}$, with $G_0 = 0$, then for $n \ge 1$,

$$|\Psi(\mathbf{G})_n(x)| \le \|\mathbf{G}\|_{\mathbf{f}} \sum_{j=1}^n |f_{n-j}(x)\,\Delta(j,j)(x)|.$$

We apply Lemma 3.2 and Lemma 3.1, (3.5), we see that

$$\|\Psi(\mathbf{G})\|_{\mathbf{f}} \le L\lambda\,\|\mathbf{G}\|_{\mathbf{f}}.$$

We therefore conclude that for small enough $\lambda > 0$ the iterates $\Psi^n(\mathbf{E})$ converge to a a fixed point which we know has to be the sequencs $\mathbf{A}$, which therefore satisfies $\|\mathbf{A} - \mathbf{E}\|_{\mathbf{f}} \le L\lambda$. So, we have proved the theorem. $\blacksquare$

3 Technical Lemmas

We recall some properties of the semigroup $\{\phi_t\}$. Of course, $\phi_t(x) = t^{-d/2}\phi\left(x/\sqrt{t}\right)$. We often write $\dot{\phi}_t$ for the derivative in t, and we write $\partial_i\phi_t$ for the partial derivatives in x_i, and $\partial_{ij}^2\phi_t$ for the second partial derivatives, etc. We also write $\Delta\phi_t \overset{\text{def}}{=} \sum_{i=1}^d \partial_{ii}^2\phi_t$, as usual. We will often use the heat equation $\dot{\phi}_t = \frac{1}{2}\Delta\phi_t$. The partial derivatives in x of ϕ are all of the form $p\phi$ for a polynomial in x whose exact form is of no concern for us. Here are some elementary properties we will use:

- If $t \le s \le 2t$ then
$$\phi_t \le 2^{d/2}\phi_s. \tag{3.1}$$
- If p is any polynomial in x, then for any $\varepsilon > 0$, there exists $C_{\varepsilon,p} > 0$ such that
$$|p(x)|\,\phi(x) \le C_{\varepsilon,p}\phi_{1+\varepsilon}(x) \tag{3.2}$$
implying
$$\left|p\left(x/\sqrt{t}\right)\right|\phi_t(x) \le C_{\varepsilon,p}\phi_{t(1+\varepsilon)}(x). \tag{3.3}$$
From this, see that derivatives in x of k-th order of $\phi_t(x)$ are bounded by $C_{\varepsilon,k}t^{-k/2}$ $\phi_{t(1+\varepsilon)}(x)$, and derivatives in t of k-th order are bounded by $C_{\varepsilon,k}t^{-k}\phi_{t(1+\varepsilon)}(x)$.

Lemma 3.1
Assume Condition 2.1, and let $\Delta(k,j)$ *be defined by (2.19). Then*

$$\sum_{j=1}^{n} |\Delta(n,j)| \le L\lambda f_n, \tag{3.4}$$

with f_n *given by (2.20)*

$$|\Delta(j,j)| \le L\lambda \left\{ \sum_{s=0}^{[j/2]} \psi_s \Gamma_{j-s} + \frac{\overline{\zeta}(j)}{j} \psi_j \right\} \tag{3.5}$$

Proof. Using (2.12) and (2.13), we can rewrite $\Delta(k,j)$ as

$$\begin{aligned}\Delta(k,j) &= \mu^{-1} a_{j-1} \left(\phi_{k\delta} - \phi_{(k-1)\delta+1}\right) - \lambda \sum_{m=1}^{j} a_m a_{j-m} \left(B_m \phi_{(k-m)\delta} - b_m \phi_{k\delta}\right) \\ &= \Delta_1(k,j) + \Delta_2(k,j),\end{aligned}$$

where

$$\Delta_2(k,j) \overset{\text{def}}{=} -\lambda \sum_{m=[k/2]+1}^{j} a_m a_{j-m} \left(B_m \phi_{(k-m)\delta} - b_m \phi_{k\delta}\right).$$

We use the convention $\sum_{m=a}^{b} = 0$ if $b < a$. As $\{a_m\}$ is bounded, we can estimate

$$|\Delta_2(k,j)| \le L\lambda \left[\sum_{s=k-j}^{[k/2]} \phi_{s\delta} \Gamma_{k-s} + \phi_{k\delta} \sum_{m=[k/2]+1}^{j} \gamma_m \right].$$

From that we see that (3.4) and (3.5) hold for Δ_2 instead of Δ, and so it remains to check these inequalities for Δ_1 :

$$\begin{aligned}\Delta_1(k,j) = \alpha &\left[\mu^{-1} \left(\phi_{k\delta} - \phi_{(k-1)\delta+1}\right) - \lambda \sum_{m=1}^{j \wedge [k/2]} a_m \left(B_m \phi_{(k-m)\delta} - b_m \phi_{k\delta}\right) \right] \\ &+ \mu^{-1} (a_{j-1} - \alpha) \left(\phi_{k\delta} - \phi_{(k-1)\delta+1}\right) \\ &- \lambda \sum_{m=1}^{j \wedge [k/2]} a_m (a_{j-m} - \alpha) \left(B_m \phi_{(k-m)\delta} - b_m \phi_{k\delta}\right), \\ &= X_1(k,j) + X_2(k,j) - X_3(k,j), \text{ say.}\end{aligned}$$

To estimate X_1, we use a Taylor approximation:

$$B_m \phi_{(k-m)\delta} = b_m \phi_{(k-m)\delta} + \overline{b}_m \dot{\phi}_{(k-m)\delta} + \frac{1}{24} E_\theta \left(\int \phi_{(k-m)\delta}^{(4)} (x - \theta y) \left[y^4\right] B_m(y)\, dy \right),$$

where E_θ refers to an expectation under the probability measure with density $4(1-\theta)^3$ on $[0,1]$. $\phi^{(4)}(z)\left[y^4\right]$ is the fourth derivative of ϕ at z in the direction y. For the error term, we use (2.4) (for $m \leq k/2$): The third summand on the right hand side is

$$\begin{aligned}
&\leq Lk^{-2}E_\theta \int \phi_{(k-m)\delta(1+\varepsilon)}(x-\theta y)\,|y|^4\,\Gamma_m(y)\,dy \\
&= Lk^{-2}E_\theta\theta^{-d} \int \phi_{(k-m)\delta(1+\varepsilon)/\theta^2}\left(\frac{x}{\theta}-y\right)|y|^4\,\Gamma_m(y)\,dy \\
&\leq Lk^{-2}\gamma^{(2)}(m)\,E_\theta\theta^{-d}\phi_{(k-m)\delta(1+\varepsilon)/\theta^2+m}\left(\frac{x}{\theta}\right) \\
&= Lk^{-2}\gamma^{(2)}(m)\,E_\theta\phi_{(k-m)\delta(1+\varepsilon)+m\theta^2}(x) \\
&\leq Lk^{-2}\gamma^{(2)}(m)\,\psi_k
\end{aligned}$$

as $\theta^2 \leq \delta(1+\varepsilon)$ if λ is small enough (by (2.17)). Furthermore

$$\begin{aligned}
\overline{b}_m\dot{\phi}_{(k-m)\delta} &= \overline{b}_m\dot{\phi}_{k\delta} + O\left(\gamma^{(1)}(m)\,mk^{-2}\psi_k\right), \\
b_m\phi_{(k-m)\delta} &= b_m\phi_{k\delta} - b_m m\delta\dot{\phi}_{k\delta} + O\left(\gamma^{(0)}(m)\,m^2k^{-2}\psi_k\right), \\
\phi_{(k-1)\delta+1} &= \phi_{k\delta} + (1-\delta)\,\dot{\phi}_{k\delta} + O\left(k^{-2}\lambda^2\psi_k\right).
\end{aligned}$$

So we get

$$X_1(k,j) = \left[\mu^{-1}(1-\delta) - \lambda\sum_{m=1}^{j\wedge(k/2)} a_m\left(\overline{b}_m - b_m m\delta\right)\right]\dot{\phi}_{k\delta} + O\left(\lambda k^{-2}\zeta_1(j\wedge[k/2])\,\psi_k\right).$$

The choice of δ was made such that the expression in square brackets is 0 if we extend the sum to ∞. Therefore, we get

$$|X_1(k,j)| \leq L\lambda\left\{k^{-2}\zeta_1(j\wedge[k/2]) + k^{-1}\zeta_2(j\wedge[k/2])\right\}\psi_k.$$

For X_2, we simply use $\phi_{(k-1)\delta+1} = \phi_{k\delta} + O\left(\lambda k^{-1}\psi_k\right)$, and Proposition 2.3 c) to get

$$|X_2(k,j)| \leq L\lambda k^{-1}\zeta_2(j),$$

and in a similar fashion, we get

$$|X_3(k,j)| = L\lambda k^{-1}\zeta_2(j\wedge[k/2])\,\psi_k.$$

Using these estimates for X_1, X_2, X_3, we get

$$\begin{aligned}
\sum_{j=1}^{n}|\Delta_1(n,j)| &\leq L\lambda\left\{n^{-2}\sum_{j=1}^{n}\zeta_1(j\wedge[n/2]) + n^{-1}\sum_{j=1}^{n}\zeta_2(j\wedge[n/2])\right\}\psi_n \\
&\leq L\lambda\left\{n^{-2}\sum_{j=1}^{n}\zeta_1(j) + n^{-1}\sum_{j=1}^{n}\zeta_2(j)\right\}\psi_n = L\lambda\overline{\zeta}(n)\,\psi_n,
\end{aligned}$$

i.e. the estimate (3.4) for Δ_1, and (3.5) is evident, too:

$$|\Delta_1(j,j)| \le L\lambda\left\{j^{-2}\zeta_1(j)+j^{-1}\zeta_2(j)\right\}\psi_n \le \frac{L\lambda}{j}\overline{\zeta}(j)\,\psi_j.$$

■

Lemma 3.2
Assume Condition 2.1 and let

$$\kappa(n) \overset{\text{def}}{=} \sum_{s=1}^{[n/2]}\psi_s\Gamma_{n-s}+\frac{\overline{\zeta}(n)}{n}\psi_n.$$

Then

$$\sum_{j=1}^{n}\kappa(j)\,f_{n-j} \le Lf_n.$$

Proof. Recall (2.20), write $f_1(n)$ for the the first of the two terms of f_n, $f_2(n)$ for the second, and similarly, we split $\kappa(n)=\kappa_1(n)+\kappa_2(n)$.

$$\begin{aligned}\sum_{j=1}^{n}\kappa_1(j)\,f_1(n-j) &= \sum_{j=1}^{n}\sum_{s=1}^{[j/2]}\psi_s\Gamma_{j-s}\sum_{t=1}^{[(n-j)/2]}t\psi_t\Gamma_{n-j-t}\\ &\le L\sum_{j=1}^{n}\sum_{s=1}^{[j/2]}\sum_{t=1}^{[(n-j)/2]}\chi_{n-s-t}(j-s)\,t\psi_{s+t}\Gamma_{n-s-t}\\ &\le L\sum_{r=1}^{[n/2]}\rho(r)\,\psi_r\Gamma_{n-r}\end{aligned}$$

with

$$\rho(r) \overset{\text{def}}{=} \sum_{j=1}^{n-r-1}\sum_{s=1}^{[j/2]\wedge(r-1)}(r-s)\,\chi_{n-r}(j-s) \le r\sum_{j=1}^{n-r-1}\sum_{s=1}^{[j/2]\wedge(r-1)}\chi_{n-r}(j-s).$$

If we fix $j-s=k$, then if $k\le(n-r)/2$, there are evidently only k possible pairs (j,s) satisfying the condition, and if $k\ge(n-r)/2$, this number is at most $n-r-k$. Therefore we get

$$\rho(r) \le \sum_{k=1}^{n-r-1}(k\wedge(n-r-k))\,\chi_{n-r}(k) \le L.$$

So we have proved

$$\sum_{j=1}^{n}\kappa_1(j)\,f_1(n-j) \le L\sum_{r=1}^{[n/2]}r\psi_r\Gamma_{n-r} \le Lf(n). \tag{3.6}$$

$$\sum_{j=1}^{n}\kappa_1(j)\,f_2(n-j)=\sum_{j=1}^{n}\sum_{s=1}^{[j/2]}\overline{\zeta}(n-j)\,\Gamma_{j-s}\psi_{n-j+s}. \tag{3.7}$$

For the summands with $j-s\leq[n/2]$ we have by (2.4) $\Gamma_{j-s}\psi_{n-j+s}\leq L\gamma(j-s)\,\psi_n\leq L\frac{\overline{\zeta}(j-s)}{(j-s)^2}\psi_n$ and by (2.9) $\overline{\zeta}(n-j)\leq L\overline{\zeta}(n)$, so we get for this part of the sum on the rhs

$$\leq L\overline{\zeta}(n)\,\psi_n\sum_{j=1}^{n}\sum_{s:s\leq[j/2],\ j-s\leq[n/2]}\frac{\overline{\zeta}(j-s)}{(j-s)^2}\leq L\overline{\zeta}(n)\,\psi_n.$$

For the summands with $j-s>[n/2]$, we get, by substituting k for $n-j+s$ that this part of the rhs of (3.7) is $\leq\sum_{k=1}^{[n/2]}\rho(k)\,\psi_k\Gamma_{n-k}$, where

$$\rho(k)=\sum_{s\leq(n-k)/2}\overline{\zeta}(k-s)\leq Lk,$$

so that we have proved

$$\sum_{j=1}^{n}\kappa_1(j)\,f_2(n-j)\leq Lf(n). \tag{3.8}$$

$$\sum_{j=1}^{n}\kappa_2(j)\,f_1(n-j)\leq Lf(n) \tag{3.9}$$

is proved in a similar way. Finally, it remains

$$\sum_{j=1}^{n}\kappa_2(j)\,f_2(n-j)=\psi_n\sum_{j=1}^{n}\frac{\overline{\zeta}(j)}{j}\overline{\zeta}(n-j).$$

The summation over $j\leq n/2$ is $\leq L\overline{\zeta}(n)\sum_j\overline{\zeta}(j)/j\leq L\overline{\zeta}(n)$ by (2.9), and the summation over $j>n/2$ is $\leq\left(\overline{\zeta}(n)/n\right)\sum_{j\leq n}\overline{\zeta}(j)\leq\overline{\zeta}(n)\sum_j\left(\overline{\zeta}(j)/j\right)\leq L\overline{\zeta}(n)$. Therefore

$$\sum_{j=1}^{n}\kappa_2(j)\,f_2(n-j)\leq Lf(n). \tag{3.10}$$

Combining (3.6), (3.8), (3.9), and (3.10) proves the claim. ■

4 Application to weakly self-avoiding walks. Proof of Theorem 1.1

We change our convention concerning the generic constant L: For the rest of the paper it denotes a constant which may depend only on the dimension d:

We derive Theorem 1.1 by applying the main Theorem 2.2 with

$$\Gamma_n\overset{\mathrm{def}}{=}Kn^{-d/2}\sum_{k=1}^{n}k^{1-d/2}\phi_{2k/5}. \tag{4.1}$$

where

$$K \overset{\text{def}}{=} \max\left(L_1(d), L_2(d)\right) \tag{4.2}$$

where L_1 is from (4.18) and L_2 from (4.22) below.

Lemma 4.1
If $d \geq 5$, then the sequence $\{\Gamma_n\}$ satisfies B1-B4 from Condition 2.1.

Proof. B2 and B4 are readily checked.
B1:

$$\begin{aligned}\Gamma_n\Gamma_m &= K^2(nm)^{-d/2}\sum_{k\leq n}\sum_{l\leq m}(kl)^{1-d/2}\phi_{2(k+l)/5}\\ &= K^2\left(\frac{n+m}{nm}\right)^{d/2}(n+m)^{-d/2}\sum_{t=2}^{n+m}\left(\sum_{k=1}^{t-1}(k(t-k))^{1-d/2}\right)\phi_{2t/5}\\ &\leq LK^2\left(\frac{n+m}{nm}\right)^{d/2}\Gamma_{n+m},\end{aligned}$$

which proves B1.

B3: We use the fact that $|y|^{2k}\phi_j \leq Lj^k\phi_{3j/2}$ for $j \in \mathbb{N}$ and $k = 0,1,2$. Therefore, we have for $m \leq t$

$$\begin{aligned}\int\phi_t(\cdot - y)|y|^{2k}\Gamma_m(y)\,dy &\leq Lm^{-d/2}\sum_{j=1}^{m}j^{1-d/2+k}\phi_{t+3j/5}\\ &\leq L\phi_{t+m}m^{-d/2}\sum_{j=1}^{m}j^{1-d/2+k} \leq L\gamma_m^{(k)}\phi_{t+m}.\end{aligned}$$

■

With this choice of Γ, we have $\bar{\zeta}(n) = O(r_n)$, r_n from (1.7), and therefore the bound in Theorem 2.2 is

$$\begin{aligned}&L\left[K\sum_{s=1}^{[n/2]}s\left(\phi_{s\delta(1+\varepsilon)}(n-s)^{-d/2}\sum_{k=1}^{n-s}k^{1-d/2}\phi_{2k/5}\right)(x) + r_n\phi_{n\delta(1+\varepsilon)}(x)\right]\\ &\leq L\left[Kn^{-d/2}\sum_{s=1}^{[n/2]}s\left(\phi_{s\delta(1+\varepsilon)}\sum_{k=1}^{n-s}k^{1-d/2}\phi_{2k/5}\right)(x) + r_n\phi_{n\delta(1+\varepsilon)}(x)\right]\\ &\leq L\left[Kn^{-d/2}\sum_{s=1}^{[n/2]}s\phi_{s\delta(1+\varepsilon)}(x) + r_n\phi_{n\delta(1+\varepsilon)}(x)\right],\end{aligned}$$

the last inequality provided

$$\delta(1+\varepsilon) \geq 4/5. \tag{4.3}$$

In order to prove Theorem 1.1 we have to show that the connectivity function in (1.5) satisfies the recursion in (1.6). This is done in Section 4.1. Finally, we have to show that the B_n's, defined through $\Pi_n = \lambda c_n^{\mathrm{SAW}} B_n$, are bounded from above by the Γ_n sequence in (4.1). This is the content of Section 4.2.

4.1 Definition of the Lace Functions

This section contains standard material on the lace expansion adapted to the model in continuous space.

Given an interval $I = [a, b] \subset \mathbb{Z}$ of integers with $0 \leq a \leq b$, we refer to a pair $\{s, t\}$ $(s < t)$ of elements of I as an *edge*. To abbreviate the notation, we write st for $\{s, t\}$. A set of edges is called a *graph*. A graph Γ on $[a, b]$ is said to be *connected* if both a and b are endpoints of edges in Γ and if, in addition, for any $c \in [a, b]$ there is an edge $st \in \Gamma$ such that $s < c < t$. Note that this is *not* in agreement with the usual notion of connectedness in graph theory. The set of all graphs on $[a, b]$ is denoted $\mathcal{B}[a, b]$, and the subset consisting of all connected graphs is denoted $\mathcal{G}[a, b]$. A *lace* is a minimally connected graph, that is, a connected graph for which the removal of any edge would result in a disconnected graph. The set of laces on $[a, b]$ is denoted $\mathcal{L}[a, b]$, and the set of laces on $[a, b]$ consisting of exactly N edges is denoted $\mathcal{L}^{(N)}[a, b]$.

A lace $\ell = \{s_1t_1, \ldots, s_Nt_N\}$ on $[0, n]$, with $s_1 = 0,\ t_N = n$, satisfies $s_i < t_{i-1},\ i = 2, \ldots, N$, and $t_i \leq s_{i+2},\ i = 1, \ldots, N-2$. We can describe the lace by the interdistances $m_1, \ldots, m_{2N-1}$ between the points s_i, t_i ordered increasingly, $s_1 = 0 < s_2 < t_1 \leq s_3 < t_2 \cdots$, i.e. $m_1 = s_2,\ m_2 = t_1 - s_2$, etc. Then of course $\sum_{i=1}^{2N-1} m_i = n$. We switch freely between the s_i-t_i-representation of the lace and the representation by the m_i, without special notice. The restrictions on the m_i are $m_i > 0$ for i even and $m_i \geq 0$ for i odd, with the additional restriction at the boundary $m_1 > 0$ and $m_{2N-1} > 0$. (For $N = 2$, all the m_i are postive). It is costumery the visualize the laces as graphs by identifying the vertices connected by a bond. Below the example of a lace with $N = 4$.



The "basic" N-lace is the graph

$$\ell_N^0 \overset{\text{def}}{=} \{(0,2), (1,4), (3,6), \ldots, (2N-5, 2N-2), (2N-3, 2N-1)\} \tag{4.4}$$

on $\{0, \ldots, 2N-1\}$. We will write $b_i = (\underline{i}, \overline{i})$ for the i-th bond in this graph, i.e. $\underline{1} = 0$, and $\underline{i} = 2i - 3$ for $i = 1, \ldots, N$, $\overline{i} = 2i$ for $i \leq N-1$, and $\overline{N} = 2N-1$. Conversely, for $i =$

$0,\dots,2N-1$, we write $\beta(i) \in \{1,\dots,N\}$ for the unique element with $i \in \left\{\underline{\beta(i)}, \overline{\beta(i)}\right\}$, i.e. $\beta(0) = 1,\ \beta(1) = 2$, etc. With this notation, we have for a lace in $\mathcal{L}^{(N)}[a,b]$

$$s_i = \sum_{j=1}^{\underline{i}} m_j,\ t_i = \sum_{j=1}^{\overline{i}} m_j.$$

If $G = \{G_t\}_{t>0}$ is any family of function in $C_*^+(\mathbb{R}^d)$, augmented by $G_0 = \delta_0$, and $\ell \in \mathcal{L}^{(N)}[0,n]$, we write with $x_0 = 0,\ x_{2N-1} = x$

$$\Xi_\ell(G,\rho)(x) \overset{\text{def}}{=} \int dx_1 \cdots dx_{2N-2} \prod_{i=1}^{2N-1} G_{m_i}(x_i - x_{i-1}) \prod_{i=1}^{n} \mathbb{I}_\rho\left(x_{\overline{i}} - x_{\underline{i}}\right). \tag{4.5}$$

For the moment, we need G only at integer values, but the more general situation is needed below.

Given a connected graph Γ on $[a,b]$, the following prescription associates to Γ a unique lace ℓ_Γ. The lace consists of edges $s_1t_1, s_2t_2, \dots$, with $t_1,\ s_1,\ t_2,\ s_2,\ \dots$ determined (in that order) by

$$\begin{aligned} t_1 &= \max\{t : at \in \Gamma\}, & s_1 &= a, \\ t_{i+1} &= \max\{t : \exists s < t_i \text{ such that } st \in \Gamma\}, & s_{i+1} &= \min\{s : st_{i+1} \in \Gamma\}. \end{aligned}$$

Given a lace ℓ, the set of all edges $st \notin \ell$ such that $\ell_{\ell\cup\{st\}} = \ell$ is denoted $\mathcal{C}(\ell)$. Edges in $\mathcal{C}(\ell)$ are said to be *compatible* with ℓ. With this formalism, we can expand the product in (1.3), obtaining

$$K_{\lambda,\rho}[a,b](\mathbf{x}) = \sum_{\Gamma\in\mathcal{B}[a,b]} \prod_{st\in\Gamma} \left(-\lambda U_{st}^\rho(\mathbf{x})\right). \tag{4.6}$$

We also define an analogous quantity, in which the sum over graphs is restricted to connected graphs, namely,

$$J[a,b](\mathbf{x}) \overset{\text{def}}{=} \sum_{\Gamma\in\mathcal{G}[a,b]} \prod_{st\in\Gamma} \left(-\lambda U_{st}^\rho(\mathbf{x})\right). \tag{4.7}$$

Recalling (1.4), this allows us to define the *lace functions*, which are the key quantities in the lace expansion:

$$\Pi_n(x_n) \overset{\text{def}}{=} \int J[0,n](\mathbf{x})\,\Phi[0,n](\mathbf{x}) \prod_{i=1}^{n-1} dx_i \tag{4.8}$$

for any $n \geq 1$ and $x_n \in \mathbb{R}^d$. The identity (1.6) is shown in the following lemma.

Lemma 4.2
For $n \geq 1$,

$$C_n^{\mathrm{SAW}} = C_{n-1}^{\mathrm{SAW}} * \phi + \sum_{k=1}^{n} \Pi_k * C_{n-k}^{\mathrm{SAW}}.$$

Proof. It suffices to show that for each path $\mathbf{x}$ we have (suppressing $\mathbf{x}$ in the formulas):

$$K[0,n] = K[1,n] + \sum_{m=1}^{n} J[0,m]\, K[m,n]. \tag{4.9}$$

Then (1.6) is obtained after insertion of (4.9) into (1.5) followed by factorization of the integral over $\mathbf{x}$. To prove (4.9), we note from (4.6) that the contribution to $K[0,n]$ from all graphs Γ for which 0 is not in an edge is exactly $K[1,n]$. To resum the contribution from the remaining graphs, we proceed as follows. When Γ does contain an edge ending at 0, we let $m[\Gamma]$ denote the largest value of m such that the set of edges in Γ with at least one end in the interval $[0,m]$ forms a connected graph on $[0,m]$. Then resummation over graphs on $[m,n]$ gives

$$K[0,n] = K[1,n] + \sum_{m=1}^{n} \sum_{\Gamma \in \mathcal{G}[0,m]} \prod_{st \in \Gamma} (-\lambda U_{st})\, K[m,n]. \tag{4.10}$$

With (4.7) this proves (4.9). ■

We next rewrite (4.8) in a form that can be used to obtain good bounds on $\Pi_n(x)$. First, splitting the sum over $\Gamma \in \mathcal{G}\,[a,b]$ according to the number of bonds in ℓ_Γ, we get

$$J[a,b] = \textstyle\sum_{N \geq 1} J_N\,[a,b]\,,$$

$$\begin{aligned} J_N\,[a,b] &\overset{\text{def}}{=} \sum_{\ell \in \mathcal{L}^{(N)}[a,b]} \sum_{\Gamma : \ell_\Gamma = \ell} \prod_{st \in \ell} (-\lambda U_{st}) \prod_{s't' \in \Gamma \setminus \ell} (-\lambda U_{s't'}) \\ &= (-\lambda)^N \sum_{\ell \in \mathcal{L}^{(N)}[a,b]} \prod_{st \in \ell} U_{st} \prod_{s't' \in \mathcal{C}(\ell)} (1 - \lambda U_{s't'}) \\ &= (-\lambda)^N\, J^{(N)}\,[a,b]\,, \text{ say.} \end{aligned} \tag{4.11}$$

Implementing into (4.8), we get a splitting

$$\Pi_n = \textstyle\sum_{N \geq 1} (-\lambda)^N\, \Pi_n^{(N)},$$

where $\Pi_n^{(N)}$ is obtained by replacing in (4.8) $J\,[0,n]$ by $J^{(N)}\,[0,n]$. Note that the sum over N is restricted to $N < n$.

An important point is that we obtain an upper bound for $\Pi_n^{(N)}$ by dropping in (4.11) the factors $(1-\lambda U_{s't'})$ for all $s't'$ which cross an endpoint of any st bond of the lace ℓ. This gives the upper bound

$$\Pi_n^{(N)}(x) \leq \sum_{\ell \in \mathcal{L}^{(N)}[0,n]} \Xi_\ell (C,\rho)(x) \tag{4.12}$$

for $N \geq 2$, where $C = \{C_n\}$. For $N=1$, there is the slight modification from "restoring" the $0n$ bond: $\Pi_n^{(1)}(x) = \Xi_{0n}(C,\rho)(x) / (1-\lambda)$.

4.2 Bounds on the lace function

We need below a slight generalization of the notion in (4.5). Given G_t, defined for real $t>0$, we define for an additional sequence $\mathbf{t} = (t_1,\ldots,t_{2N-1})$, $\Xi_\ell(G,\rho,\mathbf{t})(x)$ by replacing m_i on the right hand side of (4.5) by $m_i + t_i$. Also, given an arbitrary sequence $\mathbf{r} = (r_1,\ldots,r_{2N-1})$ of elements in $\mathbb{N}_0$, we write

$$\xi_n^{(N)}(G,\rho,\mathbf{t},\mathbf{r})(x) \stackrel{\text{def}}{=} \sum_{\mathbf{m}\in\mathcal{L}^{(N)}[0,n],\ m_i \geq r_i} \Xi_\ell(G,\rho,\mathbf{t})(x).$$

Of course, finally we are interested only in the case where the r_i are the "natural" ones from the restriction of the laces, i.e. $r_1 = r_2 = 1$, $r_3 = 0$ (if $N \geq 3$) etc. We write $\mathbf{r}^{(0)}$ for this starting sequence. If $\mathbf{t} = \mathbf{0}$ and $\mathbf{r} = \mathbf{r}^{(0)}$, we drop these arguments in the notation. We will need the more general ones in an induction argument.

We first state a simple lemma regarding normal densities.

Lemma 4.3
If $u,v,s,t>0$, $x,y\in\mathbb{R}^d$, then

$$\int \phi_u(z)\,\phi_\nu(x-z)\,\phi_s(z)\,\phi_t(y-z)\,dz \leq L\left[\frac{u+v}{uv}\right]^{d/4}\left[\frac{s+t}{st}\right]^{d/4}\phi_{u+v}(x)\,\phi_{s+t}(y). \tag{4.13}$$

Proof. By Cauchy-Schwarz the left hand side is

$$\leq \sqrt{\int \phi_u^2(z)\,\phi_\nu^2(x-z)\,dz}\sqrt{\int \phi_s^2(z)\,\phi_t^2(y-z)\,dz},$$

which equals the rhs of (4.13) by an elementary computation. ■

Let us fix some more notation. We saw that an N-lace is nothing but a sequencs $\mathbf{m} = (m_1,\ldots,m_{2N-1})$ with $\sum_i m_i = n$, and satisfying some restrictions, like $m_1 \geq 1$, $m_2 \geq 1$, $m_3 \geq 0$. We write $\mathbf{r}^{(0)} = (1,1,0,1,0,\ldots)$ for this sequence of restrictions. For an arbitrary sequence $\mathbf{r}\in\mathbb{N}_0^{2N-1}$ with $\sum_i r_i \leq n$, we write $\mathcal{L}_{\mathbf{r}}^{(N)}[0,n]$ for the set of $\mathbf{m}$ satisfying $m_i \geq r$, $\forall i$, and $\sum_i m_i = n$. The r_i need not satisfy $r_i \geq r_i^{(0)}$.

Lemma 4.4
If $\nu > 0$, $\mathbf{m} \in \mathbb{N}_0^{2N-1}$, $t_i \ge 0$, $\mathbf{x} = (x_1, \dots, x_{N-1}) \in \left(\mathbb{R}^d\right)^{N-1}$, *let*

$$\Phi^{(\nu)}_{N,\mathbf{m},\mathbf{t}}(\mathbf{x}) \overset{\text{def}}{=} \prod_{i=1}^{2N-1} \phi_{\nu m_i + t_i}\left(x_{\beta(i)-1} - x_{\beta(i-1)-1}\right),$$

with $x_0 = 0$. *If for any* i *either* $r_i \ge 1$ *or* $t_i \ge c$, *then if* $d \ge 5$, $N \ge 3$

$$\sum_{\mathbf{m}\in\mathcal{L}^{(N)}_{\mathbf{r}}[0,n]} \int dx_1 \Phi^{(\nu)}_{N,\mathbf{m},\mathbf{t}}(\mathbf{x}) \le L(c) \sum_{\mathbf{m}'\in\mathcal{L}^{(N-1)}_{\mathbf{r}'}[0,n]} \Phi^{(\nu)}_{N-1,\mathbf{m}',\mathbf{t}'}(x_2, \dots, x_{N-1}),$$

where $\mathbf{r}' \overset{\text{def}}{=} (r_3, r_1 + r_4, r_2 + r_5, r_6, \dots, r_{2N-1})$, $\mathbf{t}' \overset{\text{def}}{=} (t_3, t_1 + t_4, t_2 + t_5, t_6, \dots, t_{2N-1})$ *which both have* $2N - 3$ *components.*

Proof. The part of $\Phi^{(\nu)}_{N,\mathbf{m},\mathbf{t}}(\mathbf{x})$ which contains x_1 is

$$\phi_{m_1\nu+t_1}(x_1)\,\phi_{m_2\nu+t_2}(x_1)\,\phi_{m_4\nu+t_4}(x_2 - x_1)\,\phi_{m_5\nu+t_5}(x_3 - x_1).$$

In case $N = 3$, we have $x_3 = x_2$. Using the previous lemma for the integration over x_1, and summing over m_1, m_2, m_4, m_5, keeping $m_1 + m_4 = m_2'$, $m_2 + m_5 = m_3'$ fixed, we get for the x_1-integration and this restricted summation of the above expression a bound

$$\le L(c)\,\phi_{m_2'\nu+t_1+t_4}(x_2)\,\phi_{m_3'\nu+t_2+t_5}(x_3).$$

We write $\mathbf{m}' = (m_1' = m_3, m_1 + m_4, m_2 + m_5, m_6, \dots, m_{2N-1})$. The restrictions on the m_i' are evidently given by $m_i' \ge r_i'$. Summing over $\mathbf{m}'$ gives the desired bound.

Here is the illustration of the "collapsing mechanism":



■

Lemma 4.5
Assume $d \ge 5$. *If for some* $\nu \in \left[\frac{19}{20}, \frac{21}{20}\right]$ *and* $m \in \mathbb{N}$, $m \ge 3$, *one has*

$$G_n(x) \le \phi_{n\nu}(x), \tag{4.14}$$

for all $n < m$, *then for* $N \ge 2$, $0 < \rho \le 1$, *we have with* $L = L(d)$, *not depending on* m, N

$$\xi_m^{(N)}(G, \rho) \le L^N \rho^{Nd} \Gamma_m,$$

Γ *given by (4.1).*

Proof. We choose $\nu' \overset{\text{def}}{=} 20\nu/19$. Remark that $\nu'' \overset{\text{def}}{=} \nu' + 1/100 < 6/5$, and therefore $2\nu''/3 < 4/5$.

The assumption (4.14) implies

$$\xi_n^{(N)}(G,\rho) \le \xi_n^{(N)}\left(\phi^{(\nu)},\rho\right), \tag{4.15}$$

where $\phi^{(\nu)} = \{\phi_{\nu t}\}$.

We first want to get rid of the $\mathbb{I}_\rho$. In $\Xi_\ell\left(\phi^{(\nu)},\rho\right)(x)$, if all the m_i are ≥ 1, we can simply use $\phi_{m\nu}(x) \le L\phi_{m\nu'}(x')$ for $|x-x'| \le \rho \le 1$ from which we easily get

$$\Xi_\ell\left(\phi^{(\nu)},\rho\right)(x) \le L^N \rho^{Nd}\,\Xi_\ell\left(\phi^{(\nu')},0\right)(x).$$

There is however a complication due to the possibility of having $m_i = 0$ in the summation. Such i have to be odd, and the possibility is not present for m_1 and m_{2N-1}. Using the fact that if $m_i = 0$ then $m_{i-1}, m_{i+1} \ge 1$, we get

$$\Xi_\ell\left(\phi^{(\nu)},\rho\right)(x) \le L^N \rho^{Nd}\,\Xi_\ell\left(\phi^{(\nu')},0,\mathbf{t}^{(0)}\right)(x) \tag{4.16}$$

for all $\ell \in \mathcal{L}^{(N)}[0,n]$ where $t_i^{(0)} = 0$ for i even and $i = 1,\ 2N-1$, and $t_i^{(0)} = 1/200$ for the other i odd. Actually, the adding of the constant $1/200$ would be necessary only if m_i in fact equals 0, but there is no harm adding it always with those i for which m_i *can* be 0. It remains to estimate

$$\xi_m^{(N)}\left(\phi^{(\nu')},0,\mathbf{t}^{(0)}\right) = \sum_{\mathbf{m}\in\mathcal{L}^{(N)}[0,n]} \int dx_1 \cdots dx_{N-2}\,\Phi^{(\nu')}_{N,\mathbf{m},\mathbf{t}^{(0)}}(\mathbf{x})$$

with $x = x_{N-1}$.

For $N = 2$, there is $t_i^{(0)} = 0$ for all $i = 1,2,3$ and no integration:

$$\begin{aligned}\xi_m^{(2)}\left(\phi^{(\nu')},0\right) &\le 6 \sum_{1\le k\le l\le j,\ k+l+j=m} \phi_{k\nu'}\phi_{l\nu'}\phi_{j\nu'} \\ &\le Lm^{-d/2}\sum_{k=1}^{\lceil m/3\rceil} k^{-d/2+1}\phi_{k\nu'} \le Lm^{-d/2}\sum_{k=1}^{[m/2]} k^{-d/2+1}\phi_{4k/5} \le L\Gamma_m.\end{aligned} \tag{4.17}$$

For $N \ge 3$, we apply Lemma 4.4. Starting with $\mathbf{r}^{(0)}$ and $\mathbf{t}^{(0)}$, we recursively define $\mathbf{r}^{(k+1)} \overset{\text{def}}{=} \mathbf{r}^{(k)\prime}$, $\mathbf{t}^{(k+1)} \overset{\text{def}}{=} \mathbf{t}^{(k)\prime}$. Applying the lemma $N-2$ times we arrive at

$$\xi_m^{(N)}\left(\phi^{(\nu')},0,\mathbf{t}^{(0)}\right)(x) \le L^{N-2} \sum_{\mathbf{m}\in\mathcal{L}^{(2)}_{\mathbf{r}^{(N-2)}}[0,m]} \Phi^{(\nu')}_{2,\mathbf{m},\mathbf{t}^{(N-2)}}(x).$$

(There is no integration left when $N = 2$). The $\hat{\tau}^{(N)} \overset{\text{def}}{=} 200\mathbf{t}^{(N-2)}$, $\hat{\mathbf{r}}^{(N)} \overset{\text{def}}{=} \mathbf{r}^{(N-2)}$ can easily be computed: $\hat{\mathbf{r}}^{(2)} = (1,1,1)$, $\hat{\mathbf{r}}^{(3)} = (0,2,2)$, $\hat{\mathbf{r}}^{(4)} = (1,1,3)$, $\hat{\tau}^{(2)} =$

$(0,0,0)$, $\hat{\tau}^{(3)} = (1,0,0)$, $\hat{\tau}^{(3)} = (1,1,0)$, and $\hat{\mathbf{r}}^{(k+3)} = \hat{\mathbf{r}}^{(k)} + (1,1,1)$, $\hat{\tau}^{(k+3)} = \hat{\tau}^{(k)} + (1,1,1)$. Therefore, the only case where an $\hat{r}_i$ can be 0 is $N = 3$. Here one estimates by a similar expression as on the right hand side of (4.17) with the only difference that summation over k starts at 0, but instead of $\phi_{k\nu'}$ one has $\phi_{k\nu'+1/200}$. However, for $k = 0$, one estimates $\phi_{1/200} \leq L\phi_{\nu'}$, giving an estimate like in (4.17) with a different L. If $N > 3$, all the $\hat{r}_i^{(N)}$ are ≥ 1, and it is easily checked that $2\hat{r}_i^{(N)} \geq \hat{\tau}_i^{(N)}$. Using that, one estimates

$$\phi_{k\nu'+\hat{t}_i^{(N-2)}} \leq L\phi_{k\nu''}$$

for $k \geq \hat{r}_i^{(N-2)}$, so one gets the same estimate as in (4.17) replacing ν' by ν''. As $\nu'' < 6/5$, the argument is the same, leading to the desired estimate. ∎

Proof of Theorem 1.1. We prove by induction on m that if λ is small enough, i.e. $\lambda \leq \lambda_0(d,\varepsilon)$, we have $|B_m| \leq \Gamma_m$, where $B_m \overset{\text{def}}{=} \Pi_m/\lambda c_m$. We keep $\varepsilon > 0$ fixed, $\varepsilon \leq 1/100$.

For $m = 1$, $\Pi_1(x) = -\lambda\phi(x)\mathbb{I}_\rho(x)$, and as $c_1 = 1 - \lambda\int_{|x|\leq\rho}\phi(x)\,dx \geq 1-\lambda$, we see that with some constant $L_1(d)$ one has

$$|B_1| \leq L_1(d)\,\phi_{2/5} \leq \Gamma_1 \tag{4.18}$$

if $\lambda \leq 1/2$, where K is from (4.2). So the base of the induction is proved.

Assume now that $|B_k| \leq \Gamma_k$ for $k < m$ and define the truncated sequence $\overline{B}_k$ by B_k for $k < m$, and 0 for $k \geq m$. This sequence defines $\{\bar{C}_n\}$ via (1.1), and then $\bar{\mu}$ and $\bar{A}_n = \bar{\mu}^{-n}\bar{C}_n$. Furthermore $\bar{\delta}$ is defined by (2.16). As $|\bar{\delta}-1| \leq L\lambda$, we have $|\bar{\delta}(1+\varepsilon)-1| \leq \frac{1}{20}$ if λ is small enough. We can apply Theorem 2.2 leading to

$$\left|\bar{A}_n - \bar{a}_n\phi_{n\bar{\delta}}\right| \leq L\lambda\left[r_n\phi_{n\bar{\delta}(1+\varepsilon)} + n^{-d/2}\sum\nolimits_{j=1}^{[n/2]} j\phi_{j\bar{\delta}(1+\varepsilon)}\right].$$

As $\sup_n|\bar{a}_n - 1| \leq L\lambda$, we have

$$\bar{C}_n \leq L\bar{\mu}^n\phi_{n\bar{\delta}(1+\varepsilon)} \tag{4.19}$$

for λ small enough. As $\bar{B}_k = B_k$ for $k < m$, we have $\bar{C}_n = C_n$ for $n < m$. Remark also that by (1.2)

$$\begin{aligned}\bar{c}_m &= c_{m-1} + \lambda\sum_{k=1}^{m-1}c_kb_kc_{m-k}\\ &= c_{m-1} + \lambda\sum_{k=1}^{m}c_kb_kc_{m-k} - \lambda c_mb_m\end{aligned}$$

With the estimate (4.19), we can bound Π_m. First,

$$\Pi_m^{(1)}(x) = \mathbb{I}_\rho(x)\int\prod_{\substack{0\leq s<t\leq m,\\ st\neq 0m}}(1-\lambda U_{st}(\mathbf{x}))\,\Phi[0,n](\mathbf{x})\prod_{i=1}^{n-1}dx_i.$$

If we bound the product inside the integral from above by dropping all bonds with $t = m$, one gets

$$\begin{aligned}\Pi_m^{(1)}(x) &\leq \mathbb{I}_\rho(x)\,(\phi * C_{m-1})(x)\\ &\leq L\bar{\mu}^{m-1}\mathbb{I}_\rho(x)\,\phi_{(m-1)\bar{\delta}(1+\varepsilon)+1}(0)\\ &\leq L\bar{\mu}^{m-1}m^{-d/2}\phi_{2/5}(x) \leq L\bar{\mu}^m\Gamma_m(x)\,.\end{aligned}$$

For $\Pi_m^{(N)}$ with $N \geq 2$, we use (4.12), (4.19), and Lemma 4.5 and obtain $\Pi_m^{(N)} \leq L^N\bar{\mu}^m\Gamma_m$, and therefore, if λ is small enough $|\Pi_m| \leq L\lambda\bar{\mu}^m\Gamma_m$ implying

$$|B_m| \leq L\frac{\bar{\mu}^m}{c_m}\Gamma_m. \tag{4.20}$$

It remains to prove $\bar{\mu}^m/c_m \leq L$. Using the fact that $\bar{b}_m = 0$ and $\bar{b}_k = b_k$ for $k < m$, we get

$$\bar{c}_m = c_m(1 - \lambda b_m)\,. \tag{4.21}$$

However, $|\bar{\mu}^m/\bar{c}_m - 1| \leq L\lambda$, and from (4.20), we have $|b_m| \leq Lm^{-d/2}\bar{\mu}^m/c_m$. Using this and (4.21) and get $|b_m| \leq L$, and from that $|\bar{\mu}^m/c_m - 1| \leq L\lambda$, so we can have $\bar{\mu}^m/c_m \leq 2$ for λ small enough. This shows that with some $L_2(d)$ one has

$$|B_m| \leq L_2(d)\,\Gamma_m. \tag{4.22}$$

■

References

[1] Bolthausen, E., and Ritzmann, Ch.: *A central limit theorem for convolution equations and weakly self-avoiding walks.* ArXiV math.PR/0103218

[2] Brydges, D.C., and Spencer, T.: *Self-avoiding walks in 5 and more dimensions.* Comm. Math. Phys. **97**, 125-148 (1985)

[3] Ritzmann, Ch.: *Strong pointwise estimates for the weakly self-avoiding walk.* Ph.D. Thesis, University of Zürich, 2001.

[4] Slade, G.: *The Lace Expansion and its Application.* Springer LN in Math. **1879**, 2006.

Luca Avena, Institute of Mathematics, University of Zürich. luca.avena@math.uzh.ch
Erwin Bolthausen, Institute of Mathematics, University of Zürich. eb@math.uzh.ch
Christine Ritzmann. christine.ritzmann@gmx.ch